



CORN ALERT



Pre- and Post-Harvest Roundup® Agricultural Herbicide Applications to Roundup Ready® Corn 2

READ THE ENTIRE ROUNDUP WEATHERMAX® OR ROUNDUP POWERMAX® HERBICIDE LABEL BEFORE PROCEEDING WITH THE FOLLOWING USE DIRECTIONS. The following information is specific to and should only be used with Roundup Ready® Corn 2 products.

Roundup Ready Corn 2 is the most widely used system in corn production. Engineered to work with Roundup® agricultural herbicides, Roundup Ready Corn 2 technology provides producers with unsurpassed weed control, crop safety and added flexibility with relation to herbicide applications. One flexibility benefit producers receive with Roundup Ready Corn 2 technology is the ability to apply Roundup agricultural herbicides postemergence to manage late-season weed flushes. The following information provides specific use guidelines for pre- and post-harvest applications of Roundup WeatherMAX or Roundup PowerMAX to Roundup Ready Corn 2 products.

Preharvest Application

For late-season weed control, a preharvest application of Roundup agricultural herbicide can be applied to Roundup Ready Corn 2 products after corn reaches physiological maturity (black layer). Grain moisture could be at 35 percent or less. Roundup WeatherMAX or Roundup PowerMAX may be applied preharvest at use rates up to 22 fl oz/A.

PRECAUTIONS: Allow a minimum of 7 days between the application of this product and harvest of corn for grain or for feeding of stover.

MAXIMUM ALLOWABLE APPLICATION QUANTITIES PER SEASON.

A preharvest application may **only** be made if the combined total of previous in-crop applications of Roundup agricultural herbicide does not exceed 44 fl oz/A. The maximum combined total amount that may be applied per year is 5.3 qts/A.

Post-Harvest Application

To control weed growth after harvest, Roundup WeatherMAX or Roundup PowerMAX may be applied post-harvest at 22-32 fl oz/A. If larger weeds are present, a higher rate within the range may be necessary for control. Tank mixtures with 2,4-D or dicamba may also be used.

RESTRICTIONS: Allow a minimum of 7 days between product application and feeding of treated vegetation. For tank-mixtures also refer to other product labels.

The following table provides pre- and post-harvest application timing and maximum application rates for Roundup agricultural herbicide use with Roundup Ready Corn 2 products.

Timing	Rate
Preharvest	Maximum preharvest application rate (after maximum kernel fill is complete and the crop is physiologically mature, until 7 days before harvest) is 22 fl oz/A .
Entire Cropping Season	Combined total per year, for all applications, may not exceed (5.3 qts/A) .
Post-Harvest	22-32 fl oz/A may be applied post-harvest. A higher rate may be needed to control large weeds.




Figure 1. Two kernels are pictured. Bottom kernel is cut open to show black layer. Top kernel shows black layer at kernel tip.

ADDITIONAL APPLICATION TIPS

THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

For Ground Application:

Use the recommended rates of Roundup agricultural herbicide in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct

nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

For Aerial Applications: Use the recommended rates of Roundup agricultural herbicide in 3 to 15 gallons of spray solution per acre. Do not exceed 22 fl oz/A. See "WEEDS CONTROLLED" section on product label. AVOID DRIFT—DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

These application guidelines are from the Roundup WeatherMAX and Roundup PowerMAX herbicide label booklet. Do not combine these instructions with any other Roundup Ready corn instructions on labeling for this or other glyphosate-containing products. See "Annual Maximum Use Rate" in the "GENERAL INFORMATION" section of the Roundup WeatherMAX or Roundup PowerMAX herbicide label booklet, for additional information.

For more information, please contact your local Monsanto representative.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready® crops contain genes that confer tolerance to glyphosate, the active ingredient in Roundup® agricultural herbicides. Roundup® agricultural herbicides will kill crops that are not tolerant to glyphosate. Growing Knowledge and Design®, Roundup®, Roundup PowerMAX®, Roundup Ready®, Roundup WeatherMAX®, and Monsanto imagine® and the vine symbol are registered trademarks of Monsanto Technology LLC. DEKALB and the Winged Ear Design® are registered trademarks of DeKalb Genetics Corporation. ©2008 Monsanto Company.

A publication of
MONSANTO
imagine™

Please visit us online at www.monsantoperformance.com